

**INFORMATION IN THE FOLLOWING TABLES MAY BE USEFUL IN ANSWERING  
THE QUESTIONS IN THIS SECTION OF THE EXAMINATION.**

## PERIODIC TABLE OF THE ELEMENTS

**DO NOT DETACH FROM BOOK.**

<b>H</b>	1.0079	<b>He</b>	4
<b>Li</b>	6.941	<b>Be</b>	9.012
<b>Na</b>	22.99	<b>Mg</b>	12
<b>K</b>	39.10	<b>Ca</b>	24.30
<b>Rb</b>	85.47	<b>Sr</b>	20
<b>Cs</b>	132.91	<b>Ba</b>	38
<b>Fr</b>	87	<b>Ra</b>	40.08
	(223)		88
			226.02

§ Not yet named

58	59	60	61	62	63	64	65	66	67	68	69	70	71
Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu
140.91	144.24	(145)	150.4	151.97	157.25	158.93	162.50	164.93	167.26	168.93	173.04	174.97	
90	91	92	93	94	95	96	97	98	99	100	101	102	103
Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr
232.04	231.04	238.03	237.05	(244)	(243)	(247)	(247)	(247)	(251)	(252)	(258)	(259)	(262)

\*Lanthanide Series

†Actinide Series

**STANDARD REDUCTION POTENTIALS IN AQUEOUS SOLUTION AT 25°C**

Half-reaction		$E^\circ(V)$	
$\text{F}_2(g) + 2e^-$	$\rightarrow$	$2\text{F}^-$	2.87
$\text{Co}^{3+} + e^-$	$\rightarrow$	$\text{Co}^{2+}$	1.82
$\text{Au}^{3+} + 3e^-$	$\rightarrow$	$\text{Au}(s)$	1.50
$\text{Cl}_2(g) + 2e^-$	$\rightarrow$	$2\text{Cl}^-$	1.36
$\text{O}_2(g) + 4\text{H}^+ + 4e^-$	$\rightarrow$	$2\text{H}_2\text{O}(l)$	1.23
$\text{Br}_2(l) + 2e^-$	$\rightarrow$	$2\text{Br}^-$	1.07
$2\text{Hg}^{2+} + 2e^-$	$\rightarrow$	$\text{Hg}_2^{2+}$	0.92
$\text{Hg}^{2+} + 2e^-$	$\rightarrow$	$\text{Hg}(l)$	0.85
$\text{Ag}^+ + e^-$	$\rightarrow$	$\text{Ag}(s)$	0.80
$\text{Hg}_2^{2+} + 2e^-$	$\rightarrow$	$2\text{Hg}(l)$	0.79
$\text{Fe}^{3+} + e^-$	$\rightarrow$	$\text{Fe}^{2+}$	0.77
$\text{I}_2(s) + 2e^-$	$\rightarrow$	$2\text{I}^-$	0.53
$\text{Cu}^+ + e^-$	$\rightarrow$	$\text{Cu}(s)$	0.52
$\text{Cu}^{2+} + 2e^-$	$\rightarrow$	$\text{Cu}(s)$	0.34
$\text{Cu}^{2+} + e^-$	$\rightarrow$	$\text{Cu}^+$	0.15
$\text{Sn}^{4+} + 2e^-$	$\rightarrow$	$\text{Sn}^{2+}$	0.15
$\text{S}(s) + 2\text{H}^+ + 2e^-$	$\rightarrow$	$\text{H}_2\text{S}(g)$	0.14
$2\text{H}^+ + 2e^-$	$\rightarrow$	$\text{H}_2(g)$	0.00
$\text{Pb}^{2+} + 2e^-$	$\rightarrow$	$\text{Pb}(s)$	-0.13
$\text{Sn}^{2+} + 2e^-$	$\rightarrow$	$\text{Sn}(s)$	-0.14
$\text{Ni}^{2+} + 2e^-$	$\rightarrow$	$\text{Ni}(s)$	-0.25
$\text{Co}^2 + 2e^-$	$\rightarrow$	$\text{Co}(s)$	-0.28
$\text{Cd}^{2+} + 2e^-$	$\rightarrow$	$\text{Cd}(s)$	-0.40
$\text{Cr}^{3+} + e^-$	$\rightarrow$	$\text{Cr}^{2+}$	-0.41
$\text{Fe}^{2+} + 2e^-$	$\rightarrow$	$\text{Fe}(s)$	-0.44
$\text{Cr}^{3+} + 3e^-$	$\rightarrow$	$\text{Cr}(s)$	-0.74
$\text{Zn}^{2+} + 2e^-$	$\rightarrow$	$\text{Zn}(s)$	-0.76
$2\text{H}_2\text{O}(l) + 2e^-$	$\rightarrow$	$\text{H}_2(g) + 2\text{OH}^-$	-0.83
$\text{Mn}^{2+} + 2e^-$	$\rightarrow$	$\text{Mn}(s)$	-1.18
$\text{Al}^{3+} + 3e^-$	$\rightarrow$	$\text{Al}(s)$	-1.66
$\text{Be}^{2+} + 2e^-$	$\rightarrow$	$\text{Be}(s)$	-1.70
$\text{Mg}^{2+} + 2e^-$	$\rightarrow$	$\text{Mg}(s)$	-2.37
$\text{Na}^+ + e^-$	$\rightarrow$	$\text{Na}(s)$	-2.71
$\text{Ca}^{2+} + 2e^-$	$\rightarrow$	$\text{Ca}(s)$	-2.87
$\text{Sr}^{2+} + 2e^-$	$\rightarrow$	$\text{Sr}(s)$	-2.89
$\text{Ba}^{2+} + 2e^-$	$\rightarrow$	$\text{Ba}(s)$	-2.90
$\text{Rb}^+ + e^-$	$\rightarrow$	$\text{Rb}(s)$	-2.92
$\text{K}^+ + e^-$	$\rightarrow$	$\text{K}(s)$	-2.92
$\text{Cs}^+ + e^-$	$\rightarrow$	$\text{Cs}(s)$	-2.92
$\text{Li}^+ + e^-$	$\rightarrow$	$\text{Li}(s)$	-3.05

## ADVANCED PLACEMENT CHEMISTRY EQUATIONS AND CONSTANTS

### ATOMIC STRUCTURE

$$E = h\nu \quad c = \lambda\nu$$

$$\lambda = \frac{h}{mv} \quad p = mv$$

$$E_n = \frac{-2.178 \times 10^{-18}}{n^2} \text{ joule}$$

### EQUILIBRIUM

$$K_a = \frac{[H^+][A^-]}{[HA]}$$

$$K_b = \frac{[OH^-][HB^+]}{[B]}$$

$$K_w = [OH^-][H^+] = 1.0 \times 10^{-14} \quad @ 25^\circ C$$

$$= K_a \times K_b$$

$$pH = -\log [H^+], \quad pOH = -\log [OH^-]$$

$$14 = pH + pOH$$

$$pH = pK_a + \log \frac{[A^-]}{[HA]}$$

$$pOH = pK_b + \log \frac{[HB^+]}{[B]}$$

$$pK_a = -\log K_a, \quad pK_b = -\log K_b$$

$$K_p = K_c(RT)^{\Delta n},$$

where  $\Delta n$  = moles product gas – moles reactant gas

### THERMOCHEMISTRY/KINETICS

$$\Delta S^0 = \sum S^0 \text{ products} - \sum S^0 \text{ reactants}$$

$$\Delta H^0 = \sum \Delta H_f^0 \text{ products} - \sum \Delta H_f^0 \text{ reactants}$$

$$\Delta G^0 = \sum \Delta G_f^0 \text{ products} - \sum \Delta G_f^0 \text{ reactants}$$

$$\Delta G^0 = \Delta H^0 - T\Delta S^0$$

$$= -RT \ln K = -2.303 RT \log K$$

$$= -nFE^0$$

$$\Delta G = \Delta G^0 + RT \ln Q = \Delta G^0 + 2.303 RT \log Q$$

$$q = mc\Delta T$$

$$C_p = \frac{\Delta H}{\Delta T}$$

$$\ln[A]_t - \ln[A]_0 = -kt$$

$$\frac{1}{[A]_t} - \frac{1}{[A]_0} = kt$$

$$\ln k = \frac{-E_a}{R} \left( \frac{1}{T} \right) + \ln A$$

$E$  = energy       $v$  = velocity  
 $\nu$  = frequency       $n$  = principal quantum number  
 $\lambda$  = wavelength       $m$  = mass  
 $p$  = momentum

Speed of light,  $c = 3.0 \times 10^8 \text{ m s}^{-1}$   
 Planck's constant,  $h = 6.63 \times 10^{-34} \text{ J s}$   
 Boltzmann's constant,  $k = 1.38 \times 10^{-23} \text{ J K}^{-1}$   
 Avogadro's number =  $6.022 \times 10^{23} \text{ mol}^{-1}$   
 Electron charge,  $e = -1.602 \times 10^{-19} \text{ C}$   
 1 electron volt per atom =  $96.5 \text{ kJ mol}^{-1}$

### EQUILIBRIUM CONSTANTS

$K_a$  (weak acid)

$K_b$  (weak base)

$K_w$  (water)

$K_p$  (gas pressure)

$K_c$  (molar concentrations)

$S^0$  = standard entropy

$H^0$  = standard enthalpy

$G^0$  = standard free energy

$E^0$  = standard reduction potential

$T$  = temperature

$n$  = moles

$m$  = mass

$q$  = heat

$c$  = specific heat capacity

$C_p$  = molar heat capacity at constant pressure

$E_a$  = activation energy

$k$  = rate constant

$A$  = frequency factor

Faraday's constant,  $F = 96,500 \text{ coulombs per mole of electrons}$

Gas constant,  $R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$   
 $= 0.0821 \text{ L} \cdot \text{atm mol}^{-1} \text{ K}^{-1}$   
 $= 8.31 \text{ volt coulomb mol}^{-1} \text{ K}^{-1}$

## GASES, LIQUIDS, AND SOLUTIONS

$$PV = nRT$$

$$\left( P + \frac{n^2 a}{V^2} \right) (V - nb) = nRT$$

$$P_A = P_{total} \times X_A, \text{ where } X_A = \frac{\text{moles A}}{\text{total moles}}$$

$$P_{total} = P_A + P_B + P_C + \dots$$

$$n = \frac{m}{M}$$

$$K = {}^\circ C + 273$$

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$D = \frac{m}{V}$$

$$u_{rms} = \sqrt{\frac{3kT}{m}} = \sqrt{\frac{3RT}{M}}$$

$$KE \text{ per molecule} = \frac{1}{2} mv^2$$

$$KE \text{ per mole} = \frac{3}{2} RT$$

$$\frac{r_1}{r_2} = \sqrt{\frac{M_2}{M_1}}$$

molarity,  $M$  = moles solute per liter solution

molarity,  $m$  = moles solute per kilogram solvent

$$\Delta T_f = iK_f \times \text{molality}$$

$$\Delta T_b = iK_b \times \text{molality}$$

$$\pi = iMRT$$

$$A = abc$$

## OXIDATION-REDUCTION; ELECTROCHEMISTRY

$$Q = \frac{[C]^c [D]^d}{[A]^a [B]^b}, \text{ where } aA + bB \rightarrow cC + dD$$

$$I = \frac{q}{t}$$

$$E_{cell} = E_{cell}^0 - \frac{RT}{nF} \ln Q = E_{cell}^0 - \frac{0.0592}{n} \log Q @ 25 {}^\circ C$$

$$\log K = \frac{nE^0}{0.0592}$$

$P$  = pressure

$V$  = volume

$T$  = temperature

$n$  = number of moles

$D$  = density

$m$  = mass

$v$  = velocity

$u_{rms}$  = root-mean-square speed

$KE$  = kinetic energy

$r$  = rate of effusion

$M$  = molar mass

$\pi$  = osmotic pressure

$i$  = van't Hoff factor

$K_f$  = molal freezing-point depression constant

$K_b$  = molal boiling-point elevation constant

$A$  = absorbance

$a$  = molar absorptivity

$b$  = path length

$c$  = concentration

$Q$  = reaction quotient

$I$  = current (amperes)

$q$  = charge (coulombs)

$t$  = time (seconds)

$E^0$  = standard reduction potential

$K$  = equilibrium constant

Gas constant,  $R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$

$= 0.0821 \text{ L} \cdot \text{atm mol}^{-1} \cdot \text{k}^{-1}$

$= 8.31 \text{ volt coulomb mol}^{-1} \text{ K}^{-1}$

Boltzmann's constant,  $k = 1.38 \times 10^{-23} \text{ J K}^{-1}$

$K_f$  for H<sub>2</sub>O = 1.86 K kg mol<sup>-1</sup>

$K_b$  for H<sub>2</sub>O = 0.512 K kg mol<sup>-1</sup>

1 atm = 760 mm Hg

= 760 torr

STP = 0.000°C and 1.000 atm

Faraday's constant,  $\mathcal{F} = 96,500 \text{ coulombs per mole of electrons}$